

Inter-relationships Between Post-TBI Sequelae

Northern CA TBI Model System of Care
Santa Clara Valley Medical Center


www.tbi-sci.org

Tamara Bushnik, PhD
PAVA, January 16, 2009




Partially supported by:

- NIDRR Grant #H133A020524
- Pharmacia Inc. (now Pfizer Inc.)



Common Sequelae after TBI

- Depression
- Sleep disorders/disturbances
- Decreased/ Increased Level of activity
- Pain
- Use of Medications
- Substance use/abuse
- Fatigue



Definition of Fatigue

- “the awareness of a decreased capacity for physical and/or mental activity due to an imbalance in the availability, utilization, and/or restoration of resources needed to perform activity”


Aaronson et al, 1999



Fatigue after TBI


- Prevalence rates
 - 50%-80% in people with TBI
 - 10%-28% in people without disability

**One of the most common sequelae
after TBI**



Fatigue after TBI


- Fatigue doesn't go away
 - In a sample of individuals with TBI living in the community
 - 68% reported fatigue 2 years post-injury
 - 73% reported fatigue 5 years post-injury



Possible Contributing Factor


- Hypopituitarism
 - non-specific symptoms of pervasive fatigue, decreases in strength, poor sense of well-being overlap with those after TBI
 - In particular, the syndrome of growth hormone (GH) deficiency, gonadal, adrenal, and thyroid dysfunction

Why the Pituitary?



Hypopituitarism after TBI


	Abnormal Level/Stimulation			
	GH	Cortisol	GT	Thyroid
6-36 mnths post Agha et al, 2004	11% (11/102)	13% (13/102)	12% (12/102)	1% (1/102)
1-5.3yrs Bondanelli et al, 2004	28% (14/50)	0% (0/50)	14% (7/50)	10% (5/50)
1 year post Agha et al, 2005	10% (5/48)	19% (9/48)	12% (6/48)	2% (1/48)
1 year post Tanriverdi et al, 2006	33% (17/51)	20% (10/51)	8% (4/51)	6% (3/51)
1 year post Klose et al, 2007	29% (11/58)	3% (2/58)	2% (1/58)	2% (1/58)




Hypopituitarism after TBI

- Positive correlation between peak GH levels and
 - Verbal learning
 - Verbal short-term memory (Popovic et al, 2004)

- Positive correlation between hypopituitarism and
 - Unfavorable body composition, sleep, energy, social isolation, overall quality of life (Klose et al, 2007)



Association Between Fatigue, Severity of Injury, Duration Since Injury, and Underlying Factors




Objective

- Examine the relationship between self-reported fatigue and the following potential factors:
 - Demographic characteristics
 - Injury characteristics
 - Sleep abnormalities
 - Affective symptomatology
 - Activity patterns and limitations
 - Substance use
 - Neuroendocrine findings



Research Questions

- Endocrine abnormalities not related to time since injury
- Endocrine abnormalities related to severity of injury
- Identify unique associations between types/levels of fatigue and underlying factors




Procedure

- Participants came to Santa Clara Valley Medical Center


- Session began between 8am and 10am

- All blood tests and questionnaires completed during the 4-hour protocol



Participants

- 119 individuals with TBI
 - at least 1 year post-injury
 - living in the community
 - 16 years of age or older
 - Able to give informed consent




Participants

- Exclude people with diseases/disorders known to produce fatigue
 - Cardiovascular/pulmonary disease, diabetes mellitus, rheumatoid arthritis, multiple sclerosis, cancer, known pituitary abnormalities, chronic fatigue syndrome, pregnancy



Measures


- Demographics
- Injury severity, duration, etiology
- Barroso Fatigue Scale
- Alcohol and substance use
- Pain VAS
- Pittsburgh Sleep Quality Index (PSQI)
- Beck Depression Inventory – II (BDI-II)
- Disability Rating Scale
- Craig Handicap Reporting and Assessment Technique (CHART)
 - Cognitive Independence, Occupation, Social Integration
- Neurobehavioral Functioning Inventory (NFI)
 - Somatic, Memory/attention difficulties, Motor impairment



Barroso Fatigue Scale

- 7 subscales: Intensity, ADLs, Socialization, General Impact, Mental Functioning, Timing, Relieving Factors, Aggravating Factors

- Contains
 - Fatigue Severity Scale (FSS)
 - Multidimensional Assessment of Fatigue (MAF) subscales: Severity, ADLs, Distress, Timing, Global Fatigue Index



Measures


- Baseline blood tests:
 - CBC
 - Fasting glucose
 - Fasting basal cortisol
 - Insulin growth factor-I
 - Thyroid (free T4, TSH)
 - Testosterone (males)

- Glucagon stimulation test to assess GH response (0.03 mg/kg im, 1 mg max)



Results

- 
- ## Demographics
-
- 78 males; 41 females
 - Average age: 40 ± 12 years (16-78)
 - Duration of injury: 9 ± 7.6 years (1-37)




Demographics

<i>Marital Status</i>	
Single	45% (53)
Married	27% (32)
Sep./Div./Wid.	29% (34)
<i>Productive Activity</i>	
Employed	50% (59)
Unemployed	37% (44)
Other	13% (16)




Injury Characteristics

<i>Etiology</i>	
MVA	63% (71)
Violence	11% (12)
Falls	13% (15)
Other	13% (14)
<i>Duration of Unconsciousness</i>	
<1 d	26% (30)
1 d - <1 wk	21% (24)
1 wk - < 2 wks	15% (14)
> 2 wks	38% (44)




Measurement Scores

- Disability Rating Scale: 2.4 ± 2.0
- BDI-II: 16.0 ± 10.9
- PSQI: 7.6 ± 4.5
- CHART
 - Cognitive Independence: 76.4 ± 20.1
 - Occupation: 62.5 ± 31.3
 - Social Integration: 82.4 ± 23.0
- NFI
 - Somatic: 51.0 ± 10.2
 - Memory: 52.3 ± 10.0
 - Motor: 49.2 ± 10.2
- GFI: 26 ± 12 ; FSS: 4.4 ± 1.8



Neuroendocrine Results


Hypothyroid	12% (14)
Low cortisol (<15 mcg/dl)	64% (76)
Low testosterone (n=78)	15% (12)
Low IGF-1	19% (23)
Growth Hormone	
Severe deficiency (<3ng/ml)	34% (39)
Moderate deficiency (3-9.9 ng/ml)	31% (36)
Normal (≥ 10 ng/ml)	35% (40)



Neuroendocrine Results

	IGF-1 Level	
	Low	Normal
Growth Hormone (n=59)		
Severe deficiency	6 (26%)	17 (74%)
Moderate deficiency	3 (19%)	13 (81%)
Normal	4 (20%)	16 (80%)

$\chi^2=0.37$; $df(2)$; $p=0.83$



Time Since Injury and Endocrine Abnormalities

	Abnormal GH score	Abnormal Cortisol	Abnormal IGF-1	Abnormal T4 level	Abnormal Testosterone	Not Menstruating
Time since injury	<i>r</i> .00	.00	.07	.00	.11	.00
	<i>N</i> 114	118	117	117	Men: 77	Women: 25

	Peak GH	Cortisol	IGF-1	T4 level	TSH	Testosterone
Time since injury	<i>r</i> -.11	-.14	-.38**	-.09	.05	-.20*
	<i>N</i> 111	118	118	117	117	Men: 77

* $p < .05$ ** $p < .01$

Duration of Unconsciousness and Endocrine Abnormalities

		Abnormal GH score	Abnormal Cortisol	Abnormal IGF-1	Abnormal T4 level	Abnormal Testosterone	Not Menstruating
Duration of Unconscious	<i>r</i>	.00	.00	.14*	.00	.00	.00
	<i>N</i>	111	115	114	115	Men: 74	Women: 25

		Peak GH	Cortisol	IGF-1	T4 level	TSH	Testosterone
Duration of Unconscious	<i>r</i>	-.09	.09	.01	-.04	.01	-.02
	<i>N</i>	108	115	115	114	114	Men: 74

* $p < .05$ ** $p < .01$

Types/Levels of Fatigue and Associated Factors - Barroso

	Intensity Beta	ADLs Beta	Social Beta	Mental Beta	General Beta
Female	.28**	.26**	.17**	.25**	.17*
BDI-II	.30**	.30**	.44**	.25**	.41**
NFI Memory	.34**	---	---	.37**	---
NFI Motor	---	.30**	.24**	---	---
NFI Somatic	---	---	---	---	.21*
PSQI	.16*	---	---	---	---
CHART Social	---	.18*	---	---	---
Anti-depressant	---	---	.15*	---	---
F	33.78**	24.23**	26.47**	25.29**	22.18**
Adjusted R ²	.57	.54	.49	.40	.45

* $p < .05$ ** $p < .01$

Types/Levels of Fatigue and Associated Factors – MAF & FSS

	Severity Beta	ADLs Beta	Distress Beta	Timing Beta	GFI Beta	FSS Beta
Female	.18*	.30**	---	.25**	.21*	---
BDI-II	---	.25**	.46**	.27**	.28**	.35**
NFI Memory	.40**	---	---	.36**	.30**	---
Pain VAS	.39**	---	.22**	---	.30**	---
NFI Motor	---	.20*	---	---	---	.33**
CHART Social	---	-.21**	---	---	---	---
PSQI	---	.20*	---	---	---	---
Anti-depressant	---	---	---	---	---	-.19*
F	38.24**	17.55**	22.01**	26.51**	41.31**	25.28**
Adjusted R ²	.49	.46	.49	.42	.60	.43


* $p < .05$ ** $p < .01$

Different Types of Fatigue?

- Intensity (Barroso) – memory and sleep
- Severity (MAF) – memory and pain


- ADLs (Barroso) – motor and social
- ADLs (MAF) – motor, social and sleep

- Mental (Barroso) – memory
- General Impact (Barroso) - somatic



Limitations

- Selection bias of sample
- Cross-sectional nature
- Self-report



Thanks to

- Jeffrey Englander, MD
- Jerry Wright
- Laura Jamison
- Ketra Toda
- Kimberly Emley
- Sue Crawford, RN
- Jackie Romero, RN