

Chronic Multisymptom Illness Among Iraq/Afghanistan-Deployed US Veterans and Their Healthcare Utilization Within the Veterans Health Administration

April F. Mohanty, PhD^{1,2}, Lisa M. McAndrew, PhD^{3,4}, Drew Helmer, MD, MS³,
Matthew H. Samore, MD^{1,2,5}, and Adi V. Gundlapalli, MD, PhD^{1,2,5}

¹Informatics, Decision-Enhancement, and Analytic Sciences Center (IDEAS 2.0), VA Salt Lake City Health Care System, Salt Lake City, UT, USA;

²Division of Epidemiology, Department of Internal Medicine, University of Utah School of Medicine, Salt Lake City, UT, USA; ³Department of Veteran Affairs, War Related Illness and Injury Study Center, New Jersey Health Care System, East Orange, NJ, USA; ⁴Department of Education and Counseling Psychology, University of Albany, Albany, NY, USA; ⁵Department of Biomedical Informatics, University of Utah School of Medicine, Salt Lake City, UT, USA.

KEY WORDS: veterans; utilization; chronic disease; chronic pain; deployment health.

J Gen Intern Med

DOI: 10.1007/s11606-018-4479-6

© Society of General Internal Medicine (This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply) 2018

INTRODUCTION

As noted in prior wars such as the 1991 Gulf War, increasing evidence indicates that, chronic multisystem medically unexplained symptoms, referred to as chronic multisymptom illness (CMI), are commonly experienced by Veterans returning from Operations Enduring Freedom, Iraqi Freedom, and New Dawn (OEF/OIF/OND)^{1, 2}. Females may be at higher risk for CMI based on evidence among civilian populations and veterans from prior conflicts^{3–5}. Little is known regarding the sex-specific CMI prevalence among OEF/OIF/OND deployed veterans or related healthcare utilization. Among OEF/OIF/OND veterans nationwide and stratified by sex, we sought to estimate the prevalence of CMI-related diagnoses (hereafter referred to as CMI) and to explore trends in CMI-related (versus non-CMI-related) outpatient healthcare utilization in the Veterans Health Administration (VHA).

METHODS

Our serial cross-sectional study included 501,996 males and 69,611 females from the OEF/OIF/OND Roster provided by the Department of Defense who had encounters in any VHA facility nationwide after their last deployment end date from fiscal years 2002–2011⁶. We

defined CMI as one or more International Classification of Diseases, 9th edition, Clinical Modification (ICD-9-CM) coded diagnosis of chronic fatigue syndrome (CFS, 780.71), fibromyalgia (FMS, 729.1), or irritable bowel syndrome (IBS, 564.1). We estimated adjusted prevalence ratios (PR) for demographic and military service characteristics associated with CMI by fitting generalized linear “Log Poisson” regression models, and robust standard errors (STATA v 14.2, College Station Tx), with two-sided *p* values <0.05. We identified the settings of outpatient encounters including primary care and specialty care clinics such as gastrointestinal, pain, and rheumatology/arthritis. We defined CMI-related outpatient utilization as VHA encounters with a documented ICD-9-CM code for CMI and calculated the yearly mean number. This study was approved by our Institutional Review Board.

RESULTS

The prevalence of CMI was 4.2% in males and 8.4% in females. The higher prevalence in females versus males was consistent for CFS (3.3% versus 1.7%), FMS (9.2% versus 2.6%), and IBS (3.5% versus 1.6%). Age ≥ 30 years (PR 1.14, 95% CI 1.11–1.17) and being married (PR 1.13, 95% CI 1.10–1.16) were associated with CMI (Table 1). Enlisted rank (PR 1.35, 95% CI 1.28–1.42) and Army military branch (PR: 1.22, 95% CI 1.18–1.25) were also associated with CMI in our fully adjusted models. Over the 10-year period, the mean yearly average number of CMI-related outpatient encounters was 11.3 in females and 9.6 in males; visit counts/veteran were higher among females across all years and care settings (Fig. 1a). Total outpatient utilization was higher among veterans with CMI compared

Table 1 Demographic and Military Service Characteristics Associated with Chronic Multisymptom Illness-Related Diagnoses

Demographic characteristics	CMI dx/No CMI dx N = 27,037/544,570	Prevalence ratio (95% confidence interval)		
		Model 1	Model 2	Model 3
Male sex	5835/63,776	1 (ref)	1 (ref)	1 (ref)
Female sex	21,202/480,794	1.98 (1.93, 2.04)	2.02 (1.96, 2.07)	2.05 (1.99, 2.11)
Age at first VHA visit				
18–24 years	6705/149,455	1 (ref)	1 (ref)	1 (ref)
25–29 years	6134/130,649	1.04 (1.01, 1.08)	1.03 (1.00, 1.07)	1.02 (0.98, 1.05)
30–39 years	7829/139,540	1.24 (1.20, 1.28)	1.24 (1.20, 1.28)	1.18 (1.13, 1.22)
40+ years	6364/124,856	1.13 (1.09, 1.17)	1.16 (1.12, 1.20)	1.10 (1.06, 1.15)
Race/ethnicity				
White	12,713/272,580	1 (ref)	1 (ref)	1 (ref)
Black	2877/59,153	1.04 (1.00, 1.08)	0.91 (0.88, 0.95)	0.90 (0.87, 0.94)
Hispanic	3042/57,849	1.12 (1.08, 1.17)	1.09 (1.05, 1.14)	1.11 (1.06, 1.15)
Other	1301/26,877	1.04 (0.98, 1.10)	0.97 (0.91, 1.02)	1.02 (0.96, 1.08)
Missing	7104/128,111	1.18 (1.15, 1.21)	1.13 (1.10, 1.17)	1.21 (1.18, 1.25)
Marital status				
Not currently married	14,374/307,192	1 (ref)	1 (ref)	1 (ref)
Married	12,647/237,106	1.13 (1.11, 1.16)	1.14 (1.11, 1.17)	1.13 (1.10, 1.16)
Education				
High school or less	20,899/425,082	1 (ref)	1 (ref)	1 (ref)
Greater than high school	5791/112,449	1.05 (1.02, 1.08)	0.95 (0.92, 0.98)	1.02 (0.98, 1.05)
Military service characteristics				
Component				
Active duty	15,265/315,391	1 (ref)	1 (ref)	1 (ref)
National Guard/reserve	11,772/229,171	1.06 (1.03, 1.08)	1.09 (1.07, 1.12)	1.02 (1.00, 1.05)
Rank				
Enlisted	25,256/501,990	1 (ref)	1 (ref)	1 (ref)
Officer	1527/37,355	0.82 (0.78, 0.86)	0.74 (0.70, 0.78)	0.72 (0.68, 0.77)
Warrant	254/5217	0.97 (0.86, 1.09)	0.92 (0.81, 1.04)	0.87 (0.77, 0.98)
Branch of service				
Army	17,840/335,547	1 (ref)	1 (ref)	1 (ref)
Airforce	3175/60,547	0.99 (0.95, 1.02)	0.90 (0.87, 0.93)	0.90 (0.87, 0.94)
Navy	2953/70,546	0.80 (0.77, 0.83)	0.73 (0.70, 0.76)	0.74 (0.71, 0.77)
Marines	3048/77,309	0.75 (0.72, 0.78)	0.83 (0.80, 0.86)	0.84 (0.81, 0.87)
Coast guard	21/621	0.65 (0.43, 0.99)	0.64 (0.42, 0.97)	0.31 (0.12, 0.81)

Model 1 unadjusted

Model 2 includes age, sex, and race/ethnicity

Model 3 includes age, sex, race/ethnicity, marital status, education and military characteristics

CMI chronic multisymptom illness

to those without CMI (13.2 versus 7.1 yearly mean visits), regardless of sex (Fig. 1b). Among veterans with CMI, total outpatient utilization was higher in females versus males (14.3 versus 12.9 yearly mean visits). Among OEF/OIF/OND veterans without CMI, total outpatient utilization was higher among females versus males (7.7 versus 7.0 yearly mean visits).

DISCUSSION

Among 571,607 OEF/OIF/OND veterans who accessed VHA services from fiscal years 2002–2011, 4.7% veterans had CMI. Results from our cross-sectional study of demographic and military characteristics associated with CMI were generally consistent with prior research, including prospective studies, and extend these findings to one of the largest nationwide populations of OEF/OIF/OND veterans studied to date^{2, 3}. The finding that females have a higher prevalence of CMI is also supported by other

studies^{4, 5}. A secondary analysis of yearly prevalence of CMI, stratified by gender, showed that the prevalence increased over the 10-year period (data not shown). The extent to which more frequent healthcare utilization by females versus males overall, as observed in our study, provides more opportunities for females to receive a CMI diagnosis needs additional investigation. The finding that VHA outpatient care utilization was nearly twice as high for OEF/OIF/OND veterans with versus without CMI underscores the significant burden related to CMI both to patients and the healthcare system. Further work is needed to understand the timing or incidence and severity of symptoms, patterns of healthcare utilization within and outside the VHA, and quality of care received by male and female OEF/OIF/OND veterans with CMI. Such work may help inform updated patient-centered evidence-based guidelines and identify opportunities to improve the care of deployed veterans with CMI, a phenomenon that is being noted with regularity after major military deployments.




Fig. 1 Veterans Health Administration outpatient care utilization by OEF/OIF/OND veterans after their last deployment end date, fiscal years 2002–2011. **a** CMI-related visits/veteran by setting and sex. **b** All outpatient visits/veteran by CMI-related diagnosis status and sex. **a** CMI-related visits defined as VHA outpatient encounters associated with one or more ICD-9-CM code of chronic fatigue (780.71), fibromyalgia (729.1), or irritable bowel (564.1) syndrome. Primary care visits defined by primary stop codes: primary care (342, 348, 350, and 323) and women’s primary care (322, 339, 404, 525, 704). Specialty care visits defined by primary stop codes: gastrointestinal (307, 321, 337); pain clinic (420); rheumatology/arthritis (314). All CMI-related encounters are counted in the numerator, including those occurring on the same day. Denominator includes veterans with ≥ 1 CMI-related ICD-9-CM code in the corresponding fiscal year. **b** Outpatient encounters counted as no more than one/day. Denominator includes all veterans with ≥ 1 VHA outpatient encounter during the corresponding fiscal year. CMI, chronic multisymptom illness; ICD-9-CM, International Classification of Diseases, 9th revision, Clinical Modification; OEF, Operation Enduring Freedom; OIF, Operation Iraqi Freedom; OND, Operation New Dawn; VHA, Veterans Health Administration.

Contributors: Resources and administrative support were provided by VA Salt Lake City Health Care System (IDEAS 2.0 Center). The authors would like to acknowledge our research team members and the Veterans Informatics and Computing Infrastructure (VINCI) team in Salt Lake City.

Corresponding Author: April F. Mohanty, PhD; Informatics, Decision-Enhancement, and Analytic Sciences Center (IDEAS 2.0)VA Salt Lake City Health Care System, Salt Lake City, UT, USA (e-mail: april.mohanty@utah.edu).

Funders This work was supported by the U.S. Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development, Health Services Research and Development Project No. HIR 10-001 (PI: Samore). A.F.M. was supported by the VA Advanced Fellowship Program in Medical Informatics of the Office of Academic Affiliations, Department of Veterans Affairs, and a VA CDA 1 IK2 RX002324-01A1; and L.M.M. was supported by CDA-13-017. The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs or the US government.

Compliance with Ethical Standards:

Prior Presentation: Chronic Multisymptom Illness among Iraq/Afghanistan-Deployed Veterans and their Healthcare Utilization. Mohanty AF, McAndrew LM, Helmer D, Samore MH, Gundlapalli A. (Veterans Affairs 2017 HSR&D/QUERI National Annual Conference, Crystal City, VA, July 2017).

Conflict of Interest: The authors declare that they have no conflict of interest.

REFERENCES

1. McAndrew LM, Phillips LA, Helmer DA, Maestro K, Engel CC, Greenberg LM, et al. High healthcare utilization near the onset of medically unexplained symptoms. *J Psychosom Res.* 2017;98:98–105. <https://doi.org/10.1016/j.jpsychores.2017.05.001>
2. Smith TC, Powell TM, Jacobson IG, Smith B, Hooper TI, Boyko EJ, et al. Chronic multisymptom illness: a comparison of Iraq and Afghanistan deployers with veterans of the 1991 Gulf War. *American journal of epidemiology.* 2014;180(12):1176–87. <https://doi.org/10.1093/aje/kwu240>
3. VA/DoD Clinical Practice Guideline for the Management of Chronic Multisymptom Illness. Version 2.0 ed: Department of Veterans Affairs and the Department of Defense; 2014.
4. Vincent A, Lahr BD, Wolfe F, Clauw DJ, Whipple MO, Oh TH, et al. Prevalence of fibromyalgia: a population-based study in Olmsted County, Minnesota, utilizing the Rochester Epidemiology Project. *Arthritis Care Res (Hoboken).* 2013;65(5):786–92. <https://doi.org/10.1002/acr.21896>
5. Vincent A, Brimmer DJ, Whipple MO, Jones JF, Boneva R, Lahr BD, et al. Prevalence, incidence, and classification of chronic fatigue syndrome in Olmsted County, Minnesota, as estimated using the Rochester Epidemiology Project. *Mayo Clin Proc.* 2012;87(12):1145–52. <https://doi.org/10.1016/j.mayocp.2012.08.015>
6. Mohanty AF, Muthukutty A, Carter ME, Palmer MN, Judd J, Helmer D, et al. Chronic multisymptom illness among female Veterans deployed to Iraq and Afghanistan. *Med Care.* 2015;53(4 Suppl 1):S143–8. <https://doi.org/10.1097/MLR.0000000000000314>